

Participant Name: _____

SI MAT for OUD Learning Collaborative

Pre-Test for Session #3

Date: April 15, 2019

Instruction: Please complete the brief test before and after Session #3 by selecting one of the multi-choice options. To receive CME/CEU credits, you MUST complete pre and post tests.

- 1) A urine drug screen MUST be obtained before initiating buprenorphine treatment?
 - a. True
 - b. False

- 2) The most common daily dose of maintenance buprenorphine is?
 - a. 2mg/daily
 - b. 8mg/daily
 - c. 12mg/daily
 - d. 16mg/daily
 - e. 24mg/daily

- 3) On buprenorphine maintenance, which of the following is associated with an increased risk of dropout from treatment?
 - a. Cannabis use
 - b. Benzodiazepine use
 - c. Cannabis, Cocaine, Benzodiazepine, and/or Alcohol Use
 - d. Cocaine Use

- 4) Patients initially declining counseling will likely continue to do so once they are stabilized on MAT medication.
 - a. True
 - b. False

- 5) A patient has been in treatment for 3 months and has reduced their use of opioids, but has not completely stopped using opioids. Choose the false statement below.
 - a. Providers should terminate their buprenorphine treatment and refer out.
 - b. Providers should continue access to buprenorphine treatment.
 - c. Providers should have open discussions on the risks of mixing different substances.
 - d. Providers should continue counseling services.

- 6) True or False: Treatment with methadone and buprenorphine is associated with lower likelihood of overdose death compared with not taking these medications.
 - a. True
 - b. False